

How to take your



AGENERASE

(also called amprenavir)

Protease Inhibitor (PI)


What is Agenerase? (ah-GEN-er-ase)

Agenerase is a strong medicine that can reduce the amount of HIV in your body. Agenerase fights HIV best when you take it together with one or more other anti-HIV medicines.


How many pills do I take? And when do I take them?

Each day, take _____ pills every _____ hours.


Take your pills at these times EVERY DAY:



Morning



Afternoon



Evening

Important: To be sure there is always enough Agenerase in your body to fight HIV, take your pills as close to these times as possible. And be sure to take the right number of pills each time.

Special Instructions for taking Agenerase:

- If you eat when you take Agenerase, do NOT have a meal with a lot of fat, such as a meal with a lot of meat, cheese, or cream, or a meal with fried foods.



What if I forget to take my pills on time?

- If your next planned pill-taking time is four or more hours away, take the missed dose right away.
- If your next planned pill-taking time is less than four hours away, DO NOT take the missed pills. Instead, wait and take your next dose at the planned pill-taking time.

If you forget more than two doses in a week, call your doctor or nurse to talk about what to do.

What should I do if I have side effects?

Some people who take Agenerase have side effects (unwanted effects of the medicine). Possible side effects include:

- diarrhea
- feeling sick to your stomach or vomiting
- feeling “gassy”



If you have these or other side effects:

- Keep in mind that most side effects go away in few days or weeks
- Call your doctor or nurse. They may have ways to relieve your side effects.
- DO NOT take fewer pills or stop taking your pills. Talk to your doctor first.
- Try the ideas in the tip sheet, *“Coping with Side Effects”*.



Important: Some side effects of Agenerase (such as problems with your liver) can only be discovered by lab tests done by a doctor or nurse. So be sure to go to all your doctor’s visits.



What are the medicines I should NOT use now that I’m taking Agenerase?

Some medicines don’t mix well with Agenerase. Taking Agenerase when you are using these medicines could make you very sick or cause the medicines not to work.



Your doctor knows which medicines are safe to take with Agenerase and which are not. **Be sure you have told your doctor about ALL the medicines you are taking.** This includes:

- medicines prescribed to you by a doctor
- medicines you can buy without a prescription, including St. John’s Wort, Vitamin E, antacids, and rifampin
- herbal medicines
- street drugs

If you are also taking Videx (ddI), make sure you take Videx and Agenerase at least 1 hour apart.

Call your doctor or nurse if:



- you have questions about how to take Agenerase
- you are being bothered by side effects
- you feel like stopping or taking less Agenerase
- you are having trouble taking Agenerase on time
- you want to start taking ANY medicine you are not now taking.

DON'T FORGET: You can still spread the virus.

The anti-HIV medicines you are taking can help you stay healthier longer. But they do not cure HIV. To avoid passing HIV to others:
Practice safe sex. Don't share needles or syringes.

Developed by John Snow, Inc. and JSI Clinical Research. Boston, MA, 2002.

This document may be reproduced for educational purposes, provided it is used in its entirety and appropriate credit is given.

